

Acid-sensing ion channels interact with and inhibit BK K⁺ channels

Elena Yermolaieva Petroff*, Margaret P. Price*, Vladislav Snitsarev*, Huiyu Gong*, Victoria Korovkina†, Francois M. Abboud*†, and Michael J. Welsh*†‡§

Departments of *Internal Medicine and †Molecular Physiology and Biophysics and ‡Howard Hughes Medical Institute, Roy J. and Lucille A. Carver College of Medicine, University of Iowa, Iowa City, IA 52242

Contributed by Michael J. Welsh, December 27, 2007 (sent for review December 4, 2007)

Acid-sensing ion channels (ASICs) are neuronal non-voltage-gated cation channels that are activated when extracellular pH falls. They contribute to sensory function and nociception in the peripheral nervous system, and in the brain they contribute to synaptic plasticity and fear responses. Some of the physiologic consequences of disrupting ASIC genes in mice suggested that ASIC channels might modulate neuronal function by mechanisms in addition to their H⁺-evoked opening. Within ASIC channel's large extracellular domain, we identified sequence resembling that in scorpion toxins that inhibit K⁺ channels. Therefore, we tested the hypothesis that ASIC channels might inhibit K⁺ channel function by coexpressing ASIC1a and the high-conductance Ca²⁺- and voltage-activated K⁺ (BK) channel. We found that ASIC1a associated with BK channels and inhibited their current. Reducing extracellular pH disrupted the association and relieved the inhibition. BK channels, in turn, altered the kinetics of ASIC1a current. In addition to BK, ASIC1a inhibited voltage-gated Kv1.3 channels. Other ASIC channels also inhibited BK, although acidosis-dependent relief of inhibition varied. These results reveal a mechanism of ion channel interaction and reciprocal regulation. Finding that a reduced pH activated ASIC1a and relieved BK inhibition suggests that extracellular protons may enhance the activity of channels with opposing effects on membrane voltage. The wide and varied expression patterns of ASICs, BK, and related K⁺ channels suggest broad opportunities for this signaling system to alter neuronal function.

Acid-sensing ion channels (ASICs) are voltage-insensitive cation channels expressed in both central and peripheral neurons (1–4). ASICs are activated by extracellular protons, and several agents modify the response. Since their discovery in the early 1980s (5), they have been implicated in many physiologic processes, including nociception, mechanosensation, synaptic plasticity, and fear (1–4). They have also been implicated in pathological conditions including ischemic stroke (6, 7) and multiple sclerosis (8). These channels are formed from homomultimeric and heteromultimeric combinations of ASIC1a, -1b, -2a, -2b, and -3 subunits. Individual subunits contain short intracellular N and C termini, two transmembrane domains, and a large extracellular domain of ≈370 aa (including 14 conserved cysteines) that shows substantial sequence conservation across the degenerin/epithelial Na⁺ channel (DEG/ENaC) family. The recently published crystal structure of chicken ASIC1a shows that three subunits form a channel (9).

Activation by a reduced extracellular pH suggested that these Na⁺-conducting channels (ASIC1a also conducts Ca²⁺) would depolarize membrane voltage and thereby modify neuronal activity. In some cases the data are consistent with this action (10–12). However, some observations have not seemed to fit this proposition. For example, some results obtained in mice with disrupted ASIC genes do not seem consistent with predictions for channels producing depolarizing currents; ASIC gene disruptions and dominant-negative ASIC transgenics can paradoxically increase acid-mediated nociceptive behaviors and mechanoreceptor responses (10, 13, 14). Moreover, despite ASIC1a localization at postsynaptic membranes in central neurons, these

channels have not yet been shown to cause depolarization in response to synaptic activity (15, 16).

These incongruities caused us to wonder whether ASICs might modify neuronal function through another mechanism, and so we examined the amino acid sequence of ASICs. In the extracellular domain we identified a highly conserved sequence (R/K·Y/M·G·K·C) that resembled the part of scorpion α-K-toxins that blocks K⁺ channels (R/F/·G·K·C) (Fig. 1A) (17). In α-KTx toxins, the Lys side chain plugs the channel pore, the Arg interacts with residues in the outer vestibule of K⁺ channels, and the Cys forms a disulfide bond that stabilizes toxin structure (17–19). Although it seemed clear that the ASIC extracellular domain did not resemble an α-KTx toxin, the conserved sequence led us to hypothesize that ASICs might interact with and inhibit K⁺ channels. (After we performed these studies, the structure of chicken ASIC1a was reported; below we discuss the location of these residues in the crystal structure.) Along a similar line, Tavernarakis and Driscoll (20) reported that *Caenorhabditis elegans* degenerins and vertebrate ENaCs have amino acid sequences in their extracellular domains that show similarity to a Na⁺ channel toxin.

Results

ASIC1a Interacts with and Inhibits BK Channels. To test the hypothesis that ASICs inhibit K⁺ channels, we coexpressed ASIC1a with high-conductance Ca²⁺- and voltage-activated (BK, SLO1, or Maxi-K) K⁺ channels (21, 22). BK channels, composed of four α-subunits, are inhibited by iberitoxin (IbTx), charybdotoxin (CTx), and several other scorpion toxins containing the R/F/·G·K·C motif (Fig. 1A). These channels influence neuronal excitability, neurotransmitter release, hormone secretion, cochlear cell tuning, redox sensing, and smooth muscle tone (21–24).

Expressing BK in HEK293 cells produced depolarization-activated K⁺ currents, and mASIC1a generated H⁺-gated currents (Fig. 1B); the results were similar to previous reports (25, 26). However, when we coexpressed the two channels, BK current amplitude fell (Fig. 1B and C) even though the amount of BK protein on the cell surface did not (Fig. 1D). We obtained similar results in CHO cells and with hASIC1a (data not shown).

Protons induce conformational changes in ASIC extracellular domains and activate these channels (9, 27). Although extracellular acidosis had little effect on BK channels expressed alone (Figs. 1B and C and 4B and refs. 28 and 29), it has been reported

Author contributions: E.Y.P., M.P.P., V.S., F.M.A., and M.J.W. designed research; E.Y.P., M.P.P., V.S., and H.G. performed research; E.Y.P., M.P.P., H.G., and V.K. contributed new reagents/analytic tools; E.Y.P., M.P.P., V.S., F.M.A., and M.J.W. analyzed data; and E.Y.P., M.P.P., F.M.A., and M.J.W. wrote the paper.

The authors declare no conflict of interest.

Freely available online through the PNAS open access option.

§To whom correspondence should be addressed. E-mail: michael-welsh@uiowa.edu.

This article contains supporting information online at www.pnas.org/cgi/content/full/0712280105/DC1.

© 2008 by The National Academy of Sciences of the USA




Fig. 1. ASIC1a reduces BK current, and protons relieve inhibition. (A) Similarity between conserved sequence in ASIC extracellular domains and sequence in scorpion toxins that block K^+ channels. Bold letters indicate conserved residues in ASIC channels and the toxins. Red indicates positively charged conserved Arg and Lys residues. (B) BK currents at pH 7.4 and 2 min after change to pH 6 in HEK293 cells that were not transfected ($n = 4$), were transfected with mASIC1a ($n = 6$), were stably transfected with BK ($n = 8$), or were transfected with both BK and ASIC1a ($n = 14$). Proton-activated currents are shown in *Right*. Cells were kept at pH 7.4, and the bar indicates pH 6 application. Similar results were obtained with human ASIC1a (data not shown). Consistent with a previous report, HEK293 cells contained small endogenous H^+ -gated currents (45). (C) Current-voltage relationships for BK current in the absence ($n = 8$) and presence ($n = 14$) of mASIC1a. Data are mean \pm SEM. In the presence of mASIC1a, currents at pH 7.4 were different from those in cells with BK alone, and BK currents in the presence of ASIC1a at pH 6 and pH 7.4 were different at voltages over +40 mV ($P < 0.05$, repeated-measures ANOVA). (D) BK protein levels in the absence and presence of ASIC1a. HEK293T cells were transfected as indicated. Lysate (total protein) and biotinylated proteins were blotted with anti-BK and anti-actin antibodies.

to partially relieve CTx block (30). We tested extracellular acidosis and found that reducing pH to 6 doubled BK current amplitude in cells expressing both BK and ASIC1a (Fig. 1B and C). Thus, ASIC1a inhibited BK channel activity, and extracellular protons reduced the effect.

The ASIC1a Extracellular RYGKC Sequence Is Important for Interaction with BK. If the ASIC1a extracellular domain inhibits BK, then the extracellular domains of the two channels should interact. To test this prediction, we labeled cells expressing BK with DsRed and cells expressing ASIC1a with EGFP. We then mixed the cells together to test for cluster formation. In this assay using living cells, only the extracellular domains of proteins delivered to the cell surface will be accessible for interaction. In addition, the assay can detect weak interactions because clustering may result from multiple weak contacts between cells. We found clusters of red and green cells (Fig. 2A and B). Omitting either channel strikingly attenuated clustering. In addition, protons reduced clustering, just as they had reduced ASIC1a inhibition of BK channels. These data suggest that the extracellular domain of ASIC1a binds the extracellular surface of BK.

Earlier work showed that mutating the positively charged Arg and Lys residues in the pore-blocking motif of IbTx and CTx (Fig. 1A) to either neutral (Ala) or negative (Glu) amino acids greatly reduced their affinity for voltage-gated K^+ channels (31–33). Likewise, the corresponding mutations in the ASIC1a sequence decreased clustering (Fig. 2B). Adding IbTx also reduced clustering, suggesting that IbTx and ASIC1a might compete for BK binding. These results suggest that the extracellular domain of ASIC1a may physically interact with BK.

These results predicted that mutating conserved residues in the ASIC1a sequence would also have functional effects. Indeed, changing the cationic residues to Ala or Glu prevented BK current inhibition [Fig. 2C and supporting information (SI) Fig. 6]. The Cys in the R·F·-G·K·C sequence forms a disulfide bridge that stabilizes scorpion toxin structure, thereby positioning Arg and Lys to interact with BK. Mutating that Cys dramatically reduced CTx affinity for BK (19). We found that mASIC1a-C194A still inhibited BK current and pH 6 solution relieved inhibition (Fig. 2C). However, on switching back to pH 7.4, reinhibition of BK was markedly delayed (Fig. 2D and E), suggesting that loss of the Cys may have destabilized the structure around the RYGK sequence. These data identify ASIC1a residues that are key for BK inhibition.

The Interaction with BK Affects ASIC1a Current. In addition to the effect of ASIC1a on BK, we noticed that BK affected ASIC1a currents. Fig. 1B shows that BK altered the kinetics of mASIC1a H^+ -gated currents. Transfecting increasing amounts of BK cDNA reduced ASIC1a current amplitude and prolonged the time for desensitization (Fig. 3A–C). Thus, ASIC1a and BK exerted reciprocal effects on each other's function.

BK alteration of ASIC1a current raised the possibility that the two channels closely associate. To test this, we immunoprecipitated ASIC1a and found that it coprecipitated BK (Fig. 3D). BK also precipitated ASIC1a. We speculate that proximity in the plasma membrane positions ASIC1a and BK channels where they can interact.

Other ASIC and K^+ Channels Interact. Because other ASIC channels share a similar conserved sequence (Fig. 1A), we asked whether they would also inhibit BK current. Like ASIC1a (with an RYGKC sequence), hASIC3 (RMGKC) inhibited BK currents and pH 6 solution relieved inhibition (Fig. 4A). Previous studies showed that ASIC2a is less sensitive to pH reductions than ASIC1a or -3 and that acid fails to open ASIC2b (26, 34). We found that both ASIC2a and -2b (KYGKC) inhibited BK current but pH 6 solution failed to reverse the effects. However, more severe pH reductions attenuated the BK inhibition (Fig. 4B). When we changed the ASIC1a RYGK sequence to match that of ASIC2a (KYGKC), the mutant channel behaved like ASIC2a: it inhibited BK current, but pH 6 solution failed to relieve inhibition (Fig. 4A). Conversely, when ASIC2a contained an ASIC1a sequence (RYGKC), pH 6 solution reversed BK inhi-

performed some experiments in nominally Ca^{2+} -free extracellular solution. Under these conditions protons still increased BK current in cells expressing ASIC1a and BK. Similar results were obtained in experiments using EGTA instead of BAPTA. More direct evidence that Ca^{2+} was not required came from two other experiments. (i) ASIC2a inhibition of BK was partially relieved by protons even though ASIC2a does not conduct Ca^{2+} . (ii) Protons partially relieved BK inhibition by ASIC2b, which shows no H^{+} -gated current.

Although ASIC1a can conduct H^{+} , the stimulating effect of intracellular H^{+} on BK channels has been observed only in the absence of divalent cations (29), and in our experiments the intracellular solution contained 4.8 mM Mg^{2+} . In addition, dropping pH to values <6 elicited no additional relief of BK inhibition by ASIC1a (Fig. 4B).

- Bianchi L, Driscoll M (2002) Protons at the gate: DEG/ENaC ion channels help us feel and remember. *Neuron* 34:337–340.
- Kristhal O (2003) The ASICs: Signaling molecules? Modulators? *Trends Neurosci* 26:477–483.
- Wemmie JA, Price MP, Welsh MJ (2006) Acid-sensing ion channels: Advances, questions and therapeutic opportunities. *Trends Neurosci* 29:578–586.
- Diochot S, Salinas M, Baron A, Escoubas P, Lazdunski M (2007) Peptides inhibitors of acid-sensing ion channels. *Toxicon* 49:271–284.
- Kristhal OA, Pidoplichko VI (1980) A receptor for protons in the nerve cell membrane. *Neuroscience* 5:2325–2327.
- Xiong ZG, et al. (2004) Neuroprotection in ischemia: Blocking calcium-permeable acid-sensing ion channels. *Cell* 118:687–698.
- Yermolaieva O, Leonard AS, Schnizler MK, Abboud FM, Welsh MJ (2004) Extracellular acidosis increases neuronal cell calcium by activating acid-sensing ion channel 1a. *Proc Natl Acad Sci USA* 101:6752–6757.
- Bernardinelli L, et al. (2007) Association between the ACCN1 gene and multiple sclerosis in Central East Sardinia. *PLoS ONE* 2:e480.
- Jasti J, Furukawa H, Gonzales EB, Gouaux E (2007) Structure of acid-sensing ion channel 1 at 1.9 Å resolution and low pH. *Nature* 449:316–323.
- Price MP, et al. (2001) The DRASIC cation channel contributes to the detection of cutaneous touch and acid stimuli in mice. *Neuron* 32:1071–1083.
- Immke DC, McCleskey EW (2001) Lactate enhances the acid-sensing Na^{+} channel on ischemia-sensing neurons. *Nat Neurosci* 4:869–870.
- Vukicevic M, Kellenberger S (2004) Modulatory effects of acid-sensing ion channels (ASICs) on action potential generation in hippocampal neurons. *Am J Physiol* 287:C682–C690.
- Chen CC, Zimmer A, Sun WH, Hall J, Brownstein MJ (2002) A role for ASIC3 in the modulation of high-intensity pain stimuli. *Proc Natl Acad Sci USA* 99:8992–8997.
- Mogil JS, et al. (2005) Transgenic expression of a dominant-negative ASIC3 subunit leads to increased sensitivity to mechanical and inflammatory stimuli. *J Neurosci* 25:9893–9901.
- Wemmie JA, et al. (2002) The acid-activated ion channel ASIC contributes to synaptic plasticity, learning, and memory. *Neuron* 34:463–477.
- Alvarez de la Rosa D, et al. (2003) Distribution, subcellular localization and ontogeny of ASIC1 in the mammalian central nervous system. *J Physiol* 546:77–87.
- Miller C (1995) The charybdotoxin family of K^{+} channel-blocking peptides. *Neuron* 15:5–10.
- Gao YD, Garcia ML (2003) Interaction of agitoxin2, charybdotoxin, and iberiotoxin with potassium channels: Selectivity between voltage-gated and Maxi-K channels. *Proteins* 52:146–154.
- Drakopoulou E, et al. (1998) Consequence of the removal of evolutionary conserved disulfide bridges on the structure and function of charybdotoxin and evidence that particular cysteine spacings govern specific disulfide bond formation. *Biochemistry* 37:1292–1301.
- Tavernarakis N, Driscoll M (2000) Caenorhabditis elegans degenerins and vertebrate ENaC ion channels contain an extracellular domain related to venom neurotoxins. *J Neurogenet* 13:257–264.
- Orio P, Rojas P, Ferreira G, Latorre R (2002) New disguises for an old channel: MaxiK channel beta-subunits. *News Physiol Sci* 17:156–161.
- Salkoff L, Butler A, Ferreira G, Santi C, Wei A (2006) High-conductance potassium channels of the SLO family. *Nat Rev Neurosci* 7:921–931.
- Brenner R, et al. (2000) Vasoregulation by the beta1 subunit of the calcium-activated potassium channel. *Nature* 407:870–876.
- Tang XD, Santarelli LC, Heinemann SH, Hoshi T (2004) Metabolic regulation of potassium channels. *Annu Rev Physiol* 66:131–159.
- Korovkina VP, Brainard AM, Ismail P, Schmidt TJ, England SK (2004) Estradiol binding to maxi-K channels induces their down-regulation via proteasomal degradation. *J Biol Chem* 279:1217–1223.
- Benson CJ, et al. (2002) Heteromultimers of DEG/ENaC subunits form H^{+} -gated channels in mouse sensory neurons. *Proc Natl Acad Sci USA* 99:2338–2343.
- Adams CM, Snyder PM, Price MP, Welsh MJ (1998) Protons activate brain Na^{+} channel 1 by inducing a conformational change that exposes a residue associated with neurodegeneration. *J Biol Chem* 273:30204–30207.
- Church J, Baxter KA, McLarnon JG (1998) pH modulation of Ca^{2+} responses and a Ca^{2+} -dependent K^{+} channel in cultured rat hippocampal neurones. *J Physiol* 511:119–132.
- Avdonin V, Tang XD, Hoshi T (2003) Stimulatory action of internal protons on Slo1 BK channels. *Biophys J* 84:2969–2980.
- Perez-Cornejo P, Stampe P, Begenisch T (1998) Proton probing of the charybdotoxin binding site of Shaker K^{+} channels. *J Gen Physiol* 111:441–450.
- Park CS, Miller C (1992) Mapping function to structure in a channel-blocking peptide: Electrostatic mutants of charybdotoxin. *Biochemistry* 31:7749–7755.
- MacKinnon R, Miller C (1988) Mechanism of charybdotoxin block of the high-conductance, Ca^{2+} -activated K^{+} channel. *J Gen Physiol* 91:335–349.
- Mullmann TJ, Munujos P, Garcia ML, Giangiacomo KM (1999) Electrostatic mutations in iberiotoxin as a unique tool for probing the electrostatic structure of the maxi-K channel outer vestibule. *Biochemistry* 38:2395–2402.
- Lingueglia E, et al. (1997) A modulatory subunit of acid sensing ion channels in brain and dorsal root ganglion cells. *J Biol Chem* 272:29778–29783.
- Somodi S, et al. (2004) pH-dependent modulation of Kv1.3 inactivation: Role of His399. *Am J Physiol* 287:C1067–1076.
- Siesjo BK, Katsura K, Kristian T (1996) Acidosis-related damage. *Adv Neurol* 71:209–233, discussion 234–206.
- Katsura K, Siesjo BOK (1998) In *pH and Brain Function*, eds Kaila K, Ransom BR (Wiley-Liss, New York), p 563.
- Sausbier M, et al. (2004) Cerebellar ataxia and Purkinje cell dysfunction caused by Ca^{2+} -activated K^{+} channel deficiency. *Proc Natl Acad Sci USA* 101:9474–9478.
- Brenner R, et al. (2005) BK channel beta4 subunit reduces dentate gyrus excitability and protects against temporal lobe seizures. *Nat Neurosci* 8:1752–1759.
- Wang ZW, Saifee O, Nonet ML, Salkoff L (2001) SLO-1 potassium channels control quantal content of neurotransmitter release at the *C. elegans* neuromuscular junction. *Neuron* 32:867–881.
- Ruttiger L, et al. (2004) Deletion of the Ca^{2+} -activated potassium (BK) alpha-subunit but not the BKbeta1-subunit leads to progressive hearing loss. *Proc Natl Acad Sci USA* 101:12922–12927.
- Ramanathan K, Michael TH, Jiang GJ, Hiel H, Fuchs PA (1999) A molecular mechanism for electrical tuning of cochlear hair cells. *Science* 283:215–217.
- Veh RW, et al. (1995) Immunohistochemical localization of five members of the Kv1 channel subunits: Contrasting subcellular locations and neuron-specific co-localizations in rat brain. *Eur J Neurosci* 7:2189–2205.
- Excoffon KJ, Gansemer N, Traver G, Zabner J (2007) Functional effects of coxsackievirus and adenovirus receptor glycosylation on homophilic adhesion and adenoviral infection. *J Virol* 81:5573–5578.
- Gunthorpe MJ, Smith GD, Davis JB, Randall AD (2001) Characterisation of a human acid-sensing ion channel (hASIC1a) endogenously expressed in HEK293 cells. *Pflugers Arch* 442:668–674.
- Zhang X, Tang Y, Zhu MX (2001) Increased inwardly rectifying potassium currents in HEK-293 cells expressing murine transient receptor potential 4. *Biochem J* 354:717–725.

Cell Clustering Assay. We used methods similar to those previously reported (44).

ACKNOWLEDGMENTS. We thank Sarah K. England (University of Iowa) for HEK293 cells stably expressing BK, for the BK cDNA, and for critically reading the manuscript. We thank Leah Timmerman, Tami Nesselhauf, and the *In Vitro* and Cell Models Core (supported by National Heart, Lung, and Blood Institute Grants HL61234 and HL1515670; Cystic Fibrosis Foundation Grants R458-CR02 and ENGLH9850; and National Institute of Diabetes and Digestive and Kidney Diseases Grant DK54759). We thank Amanda Wunsch for technical help and Katherine Ashbourne Excoffon for advice. We thank Theresa Mayhew and Shawn Roach for help in preparing the manuscript. This work was supported in part by National Institutes of Health Grant HL014388 (to F.M.A.). M.J.W. is an Investigator of the Howard Hughes Medical Institute.